

For the help you need to support your course

# Facts & Formulae

mathcentre is a project offering students and staff free resources to support the transition from school mathematics to university mathematics in a range of disciplines.

[www.mathcentre.ac.uk](http://www.mathcentre.ac.uk)

This leaflet has been produced in conjunction with and is distributed by the Higher Education Academy Maths, Stats & OR Network.

For more copies contact the Network at [info@mathstore.ac.uk](mailto:info@mathstore.ac.uk)



## Algebra

Formula for solving a quadratic equation:  

$$x^2 + kx + k^2 = (x \pm k)(x^2 \pm kx + k^2)$$

$$(x + k)^2 = x^2 + 2kx + k^2, \quad (x - k)^2 = x^2 - 2kx + k^2$$

$$(x + k)(x - k) = x^2 - k^2$$
 If  $ax^2 + bx + c = 0$  then  $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

**Laws of Indices**  

$$a^m a^n = a^{m+n}, \quad \frac{a^m}{a^n} = a^{m-n}, \quad (a^m)^n = a^{mn}$$

$$a^0 = 1, \quad a^{-m} = \frac{1}{a^m}, \quad a^{1/n} = \sqrt[n]{a}, \quad a^{m/n} = (\sqrt[n]{a})^m$$

**Laws of Logarithms**  
 For any positive base  $b$  (with  $b \neq 1$ )  
 $\log_b A = c$  means  $A = b^c$   
 $\log_b A + \log_b B = \log_b AB, \quad \log_b A - \log_b B = \log_b \frac{A}{B}$   
 $n \log_b A = \log_b A^n, \quad \log_b 1 = 0, \quad \log_b b = 1$

**Formula for change of base:**  
 $\log_a x = \frac{\log_b x}{\log_b a}$   
 Logarithms to base  $e$ , denoted  $\log_e$  or alternatively  $\ln$  are called *natural logarithms*. The letter  $e$  stands for the exponential constant which is approximately 2.718.


**Partial fractions**  
 For proper fractions  $\frac{P(x)}{Q(x)}$  where  $P$  and  $Q$  are polynomials with the degree of  $P$  less than the degree of  $Q$ :  
 a linear factor  $ax + b$  in the denominator produces a partial fraction of the form  $\frac{A}{ax + b}$   
 repeated linear factors  $(ax + b)^2$  in the denominator produce partial fractions of the form  $\frac{A}{ax + b} + \frac{B}{(ax + b)^2}$   
 a quadratic factor  $ax^2 + bx + c$  in the denominator produces a partial fraction of the form  $\frac{Ax + B}{ax^2 + bx + c}$   
*Improper fractions* require an additional term which is a polynomial of degree  $n - d$  where  $n$  is the degree of the numerator and  $d$  is the degree of the denominator.

**Inequalities:**  
 $a > b$  means  $a$  is greater than  $b$   
 $a \geq b$  means  $a$  is greater than or equal to  $b$   
 $a < b$  means  $a$  is less than  $b$   
 $a \leq b$  means  $a$  is less than or equal to  $b$

## Trigonometry

**Degrees and radians**  
 $360^\circ = 2\pi$  radians,  $1^\circ = \frac{360}{2\pi} = \frac{180}{\pi}$  radians  
 $1$  radian =  $\frac{180}{\pi}$  degrees  $\approx 57.3^\circ$

**Trig ratios for an acute angle  $\theta$ :**



**Pythagoras' theorem**  
 $a^2 + b^2 = c^2$

**Standard triangles:**

$45^\circ$	$30^\circ$	$60^\circ$
$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$
$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$
$1$	$\frac{\sqrt{3}}{3}$	$\frac{2}{\sqrt{3}}$
$1$	$\frac{1}{\sqrt{3}}$	$\frac{2}{3}$
$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$
$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$
$1$	$\frac{1}{\sqrt{3}}$	$\frac{2}{\sqrt{3}}$
$1$	$\frac{\sqrt{3}}{3}$	$\frac{2}{3}$

**Common trigonometric identities**  
 $\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$   
 $\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$   
 $\tan(A \pm B) = \frac{\sin(A \pm B)}{\cos(A \pm B)} = \frac{\sin A \cos B \pm \cos A \sin B}{\cos A \cos B \mp \sin A \sin B}$   
 $2 \sin A \cos B = \sin(A + B) + \sin(A - B)$   
 $2 \cos A \cos B = \cos(A + B) + \cos(A - B)$   
 $2 \sin A \sin B = \cos(A - B) - \cos(A + B)$   
 $1 + \cot^2 A = \csc^2 A, \quad \tan^2 A + 1 = \sec^2 A$   
 $\cos 2A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A$   
 $\sin 2A = 2 \sin A \cos A$   
 $\sin^2 A = 1 - \cos^2 A$   
 $\sin^2 A = \frac{1 - \cos 2A}{2}, \quad \cos^2 A = \frac{1 + \cos 2A}{2}$   
 $\sin^2 A$  is the notation used for  $(\sin A)^2$ . Similarly  $\cos^2 A$  means  $(\cos A)^2$  etc. This notation is used with trigonometric and hyperbolic functions but with positive integer powers only.

## Hyperbolic functions

**Hyperbolic identities**  
 $e^x = \cosh x + \sinh x, \quad e^{-x} = \cosh x - \sinh x$   
 $\cosh^2 x - \sinh^2 x = 1$   
 $1 - \tanh^2 x = \operatorname{sech}^2 x$   
 $\cosh^2 x - 1 = \sinh^2 x$   
 $\sinh(x \pm y) = \sinh x \cosh y \pm \cosh x \sinh y$   
 $\cosh(x \pm y) = \cosh x \cosh y \pm \sinh x \sinh y$   
 $\sinh 2x = 2 \sinh x \cosh x$   
 $\cosh 2x = \cosh^2 x + \sinh^2 x$   
 $\sinh^2 x = \frac{\cosh 2x - 1}{2}$   
 $\cosh^2 x = \frac{\cosh 2x + 1}{2}$

**Inverse hyperbolic functions**  
 $\sinh^{-1} x = \ln(x + \sqrt{x^2 + 1})$  for  $x \geq 1$   
 $\sinh^{-1} x = \frac{1}{2} \ln \left( \frac{x+1}{x-1} \right)$  for  $-1 < x < 1$

**The Greek alphabet**

$\alpha$	alpha	$\iota$	iota	$\rho$	rho
$\beta$	beta	$\kappa$	kappa	$\sigma$	sigma
$\gamma$	gamma	$\lambda$	lambda	$\tau$	tau
$\delta$	delta	$\mu$	mu	$\upsilon$	upsilon
$\epsilon$	epsilon	$\nu$	nu	$\phi$	phi
$\zeta$	zeta	$\xi$	xi	$\chi$	chi
$\eta$	eta	$\omicron$	omicron	$\psi$	psi
$\theta$	theta	$\pi$	pi	$\omega$	omega

## Integration

$f(x)$	$\int f(x) dx = F(x) + c$	
$k$ , constant	$kx + c$	
$x^n, (n \neq -1)$	$\frac{x^{n+1}}{n+1} + c$	
$x^{-1} = \frac{1}{x}$	$\begin{cases} \ln x + c & x > 0 \\ \ln(-x) + c & x < 0 \end{cases}$	
$e^x$	$e^x + c$	
$\cos x$	$\sin x + c$	
$\sin x$	$-\cos x + c$	
$\tan x$	$\ln \sec x  + c$	$-\frac{\pi}{2} < x < \frac{\pi}{2}$
$\sec x$	$\ln \sec x + \tan x  + c$	$-\frac{\pi}{2} < x < \frac{\pi}{2}$
$\csc x$	$\ln \csc x - \cot x  + c$	$0 < x < \pi$
$\cot x$	$\ln \sin x  + c$	$0 < x < \pi$
$\cosh x$	$\sinh x + c$	
$\sinh x$	$\cosh x + c$	
$\tanh x$	$\ln \cosh x + c$	
$\coth x$	$\ln \sinh x + c$	$x > 0$
$\frac{1}{x^2 + a^2}$	$\frac{1}{a} \tan^{-1} \frac{x}{a} + c$	$a > 0$
$\frac{1}{x^2 - a^2}$	$\frac{1}{2a} \ln \frac{x-a}{x+a} + c$	$ x  > a > 0$
$\frac{1}{a^2 - x^2}$	$\frac{1}{2a} \ln \frac{a+x}{a-x} + c$	$ x  < a$
$\frac{1}{\sqrt{x^2 + a^2}}$	$\sinh^{-1} \frac{x}{a} + c$	$a > 0$
$\frac{1}{\sqrt{x^2 - a^2}}$	$\cosh^{-1} \frac{x}{a} + c$	$x \geq a > 0$
$\frac{1}{\sqrt{x^2 + k}}$	$\ln(x + \sqrt{x^2 + k}) + c$	
$\frac{1}{\sqrt{a^2 - x^2}}$	$\sin^{-1} \frac{x}{a} + c$	$-a \leq x \leq a$
$f(ax + b)$	$\frac{1}{a} F(ax + b) + c$	$a \neq 0$
e.g. $\cos(2x - 3)$	$\frac{1}{2} \sin(2x - 3) + c$	

**The linearity rule for integration**

$$\int (af(x) + bg(x)) dx = a \int f(x) dx + b \int g(x) dx, \quad (a, b \text{ constant})$$

**Integration by substitution**

$$\int f(u) \frac{du}{dx} dx = \int f(u) du \quad \text{and} \quad \int_a^b f(u) \frac{du}{dx} dx = \int_{u(a)}^{u(b)} f(u) du$$

**Integration by parts**

$$\int_a^b u \frac{dv}{dx} dx = [uv]_a^b - \int_a^b \frac{du}{dx} v dx$$

**Alternative form:**

$$\int_a^b f(x)g(x) dx = [f(x) \int g(x) dx]_a^b - \int_a^b \frac{df}{dx} \left\{ \int g(x) dx \right\} dx$$

## Differentiation

$y = f(x)$	$\frac{dy}{dx} = f'(x)$
$k$ , constant	0
$x^n$ , any constant $n$	$nx^{n-1}$
$e^x$	$e^x$
$\ln x = \log_e x$	$\frac{1}{x}$
$\sin x$	$\cos x$
$\cos x$	$-\sin x$
$\tan x = \frac{\sin x}{\cos x}$	$\sec^2 x$
$\operatorname{cosec} x = \frac{1}{\sin x}$	$-\operatorname{cosec} x \cot x$
$\sec x = \frac{1}{\cos x}$	$\sec x \tan x$
$\cot x = \frac{\cos x}{\sin x}$	$-\operatorname{cosec}^2 x$
$\sin^{-1} x$	$\frac{1}{\sqrt{1-x^2}}$
$\cos^{-1} x$	$\frac{-1}{\sqrt{1-x^2}}$
$\tan^{-1} x$	$\frac{1}{1+x^2}$
$\cosh x$	$\sinh x$
$\sinh x$	$\cosh x$
$\tanh x$	$\operatorname{sech}^2 x$
$\operatorname{sech} x$	$-\operatorname{sech} x \tanh x$
$\operatorname{cosech} x$	$-\operatorname{cosech} x \coth x$
$\coth x$	$-\operatorname{cosech}^2 x$
$\cosh^{-1} x$	$\frac{1}{\sqrt{x^2-1}}$
$\sinh^{-1} x$	$\frac{1}{\sqrt{x^2+1}}$
$\tanh^{-1} x$	$\frac{1}{1-x^2}$

**The linearity rule for differentiation**

$$\frac{d}{dx}(au + bv) = a \frac{du}{dx} + b \frac{dv}{dx} \quad a, b \text{ constant}$$

**The product and quotient rules for differentiation**

$$\frac{d}{dx}(uv) = u \frac{dv}{dx} + v \frac{du}{dx} \quad \frac{d}{dx} \left( \frac{u}{v} \right) = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

**The chain rule for differentiation**

If  $y = y(u)$  where  $u = u(x)$  then  $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$

For example,  
 if  $y = (\cos x)^{-1}$ ,  $\frac{dy}{dx} = -1(\cos x)^{-2}(-\sin x)$

$$\begin{bmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{bmatrix} \begin{bmatrix} j & k \\ l & m \\ n & o \end{bmatrix} = \begin{bmatrix} a_1j + a_2l + a_3n & a_1k + a_2m + a_3o \\ b_1j + b_2l + b_3n & b_1k + b_2m + b_3o \end{bmatrix}$$

If  $\mathbf{a} = a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}$  and  $\mathbf{b} = b_1\mathbf{i} + b_2\mathbf{j} + b_3\mathbf{k}$  then

$\mathbf{a} \cdot \mathbf{b} = a_1b_1 + a_2b_2 + a_3b_3$   
 If  $\mathbf{a} = a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}$  and  $\mathbf{b} = b_1\mathbf{i} + b_2\mathbf{j} + b_3\mathbf{k}$  then

$\mathbf{a} \times \mathbf{b} = (a_2b_3 - a_3b_2)\mathbf{i} + (a_3b_1 - a_1b_3)\mathbf{j} + (a_1b_2 - a_2b_1)\mathbf{k}$

If  $\mathbf{a} = a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}$  and  $\mathbf{b} = b_1\mathbf{i} + b_2\mathbf{j} + b_3\mathbf{k}$  then  
 $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}| |\mathbf{b}| \cos \theta$

If  $\mathbf{a} = a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}$  and  $\mathbf{b} = b_1\mathbf{i} + b_2\mathbf{j} + b_3\mathbf{k}$  then  
 $\mathbf{a} \times \mathbf{b} = |\mathbf{a}| |\mathbf{b}| \sin \theta \mathbf{e}$   
 where  $\mathbf{e}$  is a unit vector perpendicular to the plane containing  $\mathbf{a}$  and  $\mathbf{b}$  in a sense defined by the right hand screw rule.

### Vectors

If  $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$  then  $|\mathbf{r}| = \sqrt{x^2 + y^2 + z^2}$   
 If  $\mathbf{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$  then  $\mathbf{r}$  may be used to denote  $\sqrt{-1}$ .

Relationship between hyperbolic and trig functions  
 $\cosh^2 x - \sinh^2 x = 1$   
 $\cosh x = \frac{e^x + e^{-x}}{2}$   
 $\sinh x = \frac{e^x - e^{-x}}{2}$

Euler's relations  
 $e^{j\theta} = \cos \theta + j \sin \theta$   
 $e^{-j\theta} = \cos \theta - j \sin \theta$   
 $e^{j\theta} e^{-j\theta} = 1$   
 $e^{j\theta} = \cos \theta + j \sin \theta$   
 $e^{-j\theta} = \cos \theta - j \sin \theta$

Cartesian form:  $z = a + bj$   
 Polar form:  $z = r(\cos \theta + j \sin \theta) = r e^{j\theta}$   
 where  $\theta = \tan^{-1} \frac{b}{a}$

De Moivre's theorem  
 If  $z = r e^{j\theta}$ , then  $z^n = r^n e^{jn\theta}$   
 $z^{1/z} = r^{1/z} e^{j\theta/z}$   
 $z^{1/z} = r^{1/z} (\cos \frac{\theta}{z} + j \sin \frac{\theta}{z})$

Multiplication and division in polar form  
 $z_1 z_2 = r_1 r_2 e^{j(\theta_1 + \theta_2)}$   
 $\frac{z_1}{z_2} = \frac{r_1}{r_2} e^{j(\theta_1 - \theta_2)}$

Complex Numbers  
 Cartesian form:  $z = a + bj$   
 Polar form:  $z = r(\cos \theta + j \sin \theta) = r e^{j\theta}$   
 where  $\theta = \tan^{-1} \frac{b}{a}$   
 Euler's relations  
 $e^{j\theta} = \cos \theta + j \sin \theta$   
 $e^{-j\theta} = \cos \theta - j \sin \theta$   
 $e^{j\theta} e^{-j\theta} = 1$   
 De Moivre's theorem  
 If  $z = r e^{j\theta}$ , then  $z^n = r^n e^{jn\theta}$   
 $z^{1/z} = r^{1/z} e^{j\theta/z}$   
 $z^{1/z} = r^{1/z} (\cos \frac{\theta}{z} + j \sin \frac{\theta}{z})$

Exponential functions  
 Graph of  $y = e^x$  showing exponential growth  
 Graph of  $y = e^{-x}$  showing exponential decay

Graphs of  $y = 0.5^x$ ,  $y = 3^x$ , and  $y = 2^x$

Logarithmic functions  
 Graphs of  $y = \ln x$  and  $y = \log_{10} x$

Hyperbolic functions  
 Graphs of  $y = \sinh x$ ,  $y = \cosh x$  and  $y = \tanh x$

The sine rule and cosine rule  
 The sine rule  
 $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$   
 The cosine rule  
 $a^2 = b^2 + c^2 - 2bc \cos A$

Trigonometric functions  
 The sine and cosine functions are periodic with period  $2\pi$ .  
 The tangent function is periodic with period  $\pi$ .

Inverse trigonometric functions  
 $y = \sin^{-1} x$   
 $y = \cos^{-1} x$   
 $y = \tan^{-1} x$

Arithmetic progression:  $a, a+d, a+2d, \dots$   
 Sum of first  $n$  terms,  $S_n = \frac{n}{2}(2a + (n-1)d)$   
 kth term  $= a + (k-1)d$

Geometric progression:  $a, ar, ar^2, \dots$   
 Sum of  $n$  terms,  $S_n = \frac{a(1-r^n)}{1-r}$ , provided  $r \neq 1$   
 kth term  $= ar^{k-1}$   
 $a$  = first term,  $r$  = common ratio

Sum of the squares of the first  $n$  integers,  
 $1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{1}{6}n(n+1)(2n+1)$

Sum of an infinite geometric series:  
 $S_\infty = \frac{a}{1-r}$ ,  $-1 < r < 1$

The binomial theorem  
 If  $n$  is a positive integer  
 $(1+x)^n = 1 + nx + \frac{n(n-1)}{2}x^2 + \dots + x^n$   
 When  $n$  is negative or fractional, the series is infinite and converges when  $-1 < x < 1$

Standard power series expansions  
 $e^x = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \dots$   
 $\sin x = x - \frac{x^3}{6} + \frac{x^5}{120} - \dots$   
 $\cos x = 1 - \frac{x^2}{2} + \frac{x^4}{24} - \dots$

The exponential function as the limit of a sequence  
 $\lim_{n \rightarrow \infty} \left(1 + \frac{x}{n}\right)^n = e^x$

Pascal's triangle:  
 The pattern of the coefficients is seen in  $(1+x)^n$ , so, for example,  $4! = 1 \cdot 2 \cdot 3 \cdot 4$   
 $0! = 1, n! = n(n-1) \dots 1$

The coefficient of  $x^k$  in the binomial expansion of  $(1+x)^n$  is  $\binom{n}{k}$  or  ${}^nC_k$ , when  $n$  is a positive integer is denoted by  $\binom{n}{k}$  or  ${}^nC_k$ .  
 ${}^nC_k$  is the number of subsets with  $k$  elements that can be chosen from a set with  $n$  elements.

Remember that  $AB \neq BA$  except in special cases.

Matrix multiplication: for  $2 \times 2$  matrices  
 provided that  $ad - bc \neq 0$ .  
 If  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  then  $A^{-1} = \frac{1}{ad-bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$   
 The inverse of a  $2 \times 2$  matrix (expanded along the first row).

The  $3 \times 3$  matrix  $A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$  has determinant  
 $|A| = a_{11}(a_{22}a_{33} - a_{23}a_{32}) - a_{12}(a_{21}a_{33} - a_{23}a_{31}) + a_{13}(a_{21}a_{32} - a_{22}a_{31})$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

Graphs of common functions  
 Linear  $y = mx + c$ ,  $m$ =gradient,  $c$  = vertical intercept

The equation of a circle centre  $(a, b)$ , radius  $r$   
 $(x-a)^2 + (y-b)^2 = r^2$

Quadratic functions  $y = ax^2 + bx + c$   
 (1)  $a > 0$   
 (2)  $b^2 - 4ac < 0$   
 (3)  $b^2 - 4ac = 0$   
 (1)  $a < 0$   
 (2)  $b^2 - 4ac > 0$   
 (3)  $b^2 - 4ac = 0$

Completing the square  
 If  $a \neq 0$ ,  $ax^2 + bx + c = a\left(x + \frac{b}{2a}\right)^2 + \frac{4ac - b^2}{4a}$

The modulus function  
 $|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$

The unit step function,  $u(x)$   
 $u(x) = \begin{cases} 1 & \text{if } x \geq 0 \\ 0 & \text{if } x < 0 \end{cases}$

The sine rule  
 $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$

The cosine rule  
 $a^2 = b^2 + c^2 - 2bc \cos A$

Pascal's triangle:  
 The pattern of the coefficients is seen in  $(1+x)^n$ , so, for example,  $4! = 1 \cdot 2 \cdot 3 \cdot 4$   
 $0! = 1, n! = n(n-1) \dots 1$

The coefficient of  $x^k$  in the binomial expansion of  $(1+x)^n$  is  $\binom{n}{k}$  or  ${}^nC_k$ , when  $n$  is a positive integer is denoted by  $\binom{n}{k}$  or  ${}^nC_k$ .  
 ${}^nC_k$  is the number of subsets with  $k$  elements that can be chosen from a set with  $n$  elements.

Remember that  $AB \neq BA$  except in special cases.

Matrix multiplication: for  $2 \times 2$  matrices  
 provided that  $ad - bc \neq 0$ .  
 If  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  then  $A^{-1} = \frac{1}{ad-bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$   
 The inverse of a  $2 \times 2$  matrix (expanded along the first row).

The  $3 \times 3$  matrix  $A = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$  has determinant  
 $|A| = a_{11}(a_{22}a_{33} - a_{23}a_{32}) - a_{12}(a_{21}a_{33} - a_{23}a_{31}) + a_{13}(a_{21}a_{32} - a_{22}a_{31})$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$

The  $2 \times 2$  matrix  $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  has determinant  
 $|A| = ad - bc$